

AD-A078 774

FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS F/G 12/1
THE EXTREME POINTS OF THE SET OF DECREASING FAILURE RATE DISTRI--ETC(U)
AUG 79 N A LANGBERG , R V LEON , J LYNCH AFOSR-78-3678
UNCLASSIFIED FSU-STATISTICS-M484-REV AFOSR-TR-79-1219 NL

[OF]
AD
50/114



END
DATE
FORMED
1-80
DOC



ADA 078774

DDC FILE COPY

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER AFOSR-TR-79-1219	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitle) THE EXTREME POINTS OF THE SET OF DECREASING FAILURE RATE DISTRIBUTIONS		5. TYPE OF REPORT & PERIOD COVERED Interim	
7. AUTHOR(s) Naftali A. Langberg, Ramon V. Leon, James Lynch, and Frank Proschan		6. PERFORMING ORG. REPORT NUMBER 78-5-Revised	
9. PERFORMING ORGANIZATION NAME AND ADDRESS The Florida State University Department of Statistics Tallahassee, FL 32306		8. CONTRACT OR GRANT NUMBER(s) AFOSR-78-3678 NSF-11279-84678	
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM Bolling AFB, Washington, DC 32306		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2304/A5	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 14781-STATISTICS-11-11-11 FV		12. REPORT DATE Aug 79	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		13. NUMBER OF PAGES 16	
17. DISTRIBUTION STATEMENT (of the Abstract in Block 20, if different from Report) DDC RECEIVED DEC 27 1979 E		15. SECURITY CLASS. (of this Report) UNCLASSIFIED	
18. SUPPLEMENTARY NOTES Supersedes AD-A064786			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Decreasing failure rate, convex set, extreme points, Choquet's Theorem mixture, hazard function, failure rate, reliability. infinity 9 12 19 038			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Since the class of extended decreasing failure rate (EDFR) life distributions (i.e., distributions with support in $[0, \infty)$ is compact and convex, it follows from Choquet's Theorem that every EDFR life distribution can be represented as a mixture of extreme points of the EDFR class. We identify the extreme points of this class and of the standard class of decreasing failure rate (DFR) life distributions. Further, we show that even though the convex class of DFR life distributions is not compact, every DFR life distribution can be represented as a mixture of extreme points of the DFR class.			

THE EXTREME POINTS OF THE SET OF DECREASING
FAILURE RATE DISTRIBUTIONS

by

Naftali A. Langberg¹, Ramón V. León¹,
James Lynch² and Frank Proschan¹

FSU Statistics Report M484-Revised
AFOSR Technical Report No. 78-5-Revised

August, 1979

The Florida State University
Department of Statistics
Tallahassee, Florida 32306

¹Research sponsored by the Air Force Office of Scientific Research, AFSC, USAF, under Grant AFOSR 78-3678. Affiliation: Department of Statistics, Florida State University.

²Research sponsored by the National Science Foundation MCS-7904698. Affiliation: Department of Statistics, Pennsylvania State University.

AMS 1970 Subject Classifications: Primary 62E10, 62N05, Secondary 60K99.

The Extreme Points of the Set of Decreasing
Failure Rate Distributions

by

Naftali A. Langberg, Ramón V. León,
James Lynch and Frank Proschan

ABSTRACT.

Since the class of extended decreasing failure rate (EDFR) life distributions (i.e., distributions with support in $[0, \infty]$) is compact and convex, it follows from Choquet's Theorem that every EDFR life distribution can be represented as a mixture of extreme points of the EDFR class. We identify the extreme points of this class and of the standard class of decreasing failure rate (DFR) life distributions. Further, we show that even though the convex class of DFR life distributions is not compact, every DFR life distribution can be represented as a mixture of extreme points of the DFR class.

AMS 1970 Subject Classifications. Primary 62E10, 62N05; Secondary 60K99.

Key Words: Decreasing failure rate, convex set, extreme points, Choquet's Theorem, mixture, hazard function, failure rate, reliability.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
This technical report has been reviewed and is
approved for public release under AFR 190-12 (7b).
Distribution is unlimited.
A. D. BLOSE
Technical Information Officer

1. Introduction and Summary.

The class of decreasing failure rate (DFR) distributions plays an important role in the theory and application of reliability, biometry, and other fields of statistics. (See, e.g., Barlow and Proschan, 1975, Chaps. 3 and 4, and Proschan, 1963).

Such distributions govern the lifelengths of systems which do not age adversely over time in the sense that the conditional survival probability given the age of the system is an increasing function of the age. For example, DFR distributions govern the lifelengths (i) of metals subject to "work-hardening", (ii) of many solid state components, (iii) of businesses, (iv) of mixtures of exponential distributions, etc. .

The DFR class is convex, as is the class of extended decreasing failure rate distributions (EDFR) which contains distributions placing mass at ∞ . This latter class is also compact in the topology of weak convergence of probability measures. (See Section 2.)

From the Krein-Milman Theorem and Choquet's Theorem, we know that the basic building blocks of convex compact sets are their extreme points. In particular, it follows from Choquet's Theorem, stated in Section 2, that the set of extreme points is the smallest set of EDFR distributions with the property that every EDFR distribution may be represented as a mixture of its elements.

The main purpose of this paper is to identify the extreme points of the EDFR class (Section 3). They are those EDFR distributions having failure rate functions whose derivatives are close to zero in an appropriate sense. More specifically, an EDFR distribution F is not an extreme point of

Accession For	NTIS GAO&I	
	DEC IAB	
	Unannounced	
	Publication	
Dist		
Availability Codes		
Avail and/or		
Special		
Dist		A

the EDFR class if and only if (i) $0 < F(0) < 1$, (ii) $0 < F(\infty) < 1$, or (iii) the derivative of the failure rate function is a.e. uniformly bounded away from 0 in some interval (see Theorem 3.1).

Since the DFR class is an extremal subset of the EDFR class, it follows that a distribution F is an extreme point of the DFR class if and only if F is an extreme point of the EDFR class and it places no mass at ∞ . Thus the extreme points of the DFR class are also identified (see Corollary 4.2).

In Theorem 4.5 we show that even though the convex class of DFR distributions is not compact, the set of extreme points of this class is the smallest set of DFR distributions with the property that every DFR distribution may be represented as a mixture of its elements.

2. Preliminaries.

Let F be an extended life distribution; i.e., F is a distribution function possibly placing mass at ∞ such that $F(0-) = 0$. When an extended life distribution places no mass at ∞ , it is simply called a life distribution. The function $\bar{F} \equiv 1 - F$ is called the survival probability, and $R_F \equiv -\ln \bar{F}$ is called the hazard function. If R_F is absolutely continuous on every closed interval contained in $(0, \infty)$, then any measurable function r_F almost everywhere (a.e.) equal to R'_F , the derivative of R_F , is called a failure rate function. (Throughout, measurable means Borel measurable and all measures are Borel measures). When F has a density f , the failure rate function $r_F = f/\bar{F}$ a.e. . When the distribution F is clearly understood, the subscript f will be suppressed.

Definition 2.1. Let F be an extended life distribution. Then F is said to be an extended decreasing failure rate (EDFR) distribution if R is concave on $(0, \infty)$. If an EDFR distribution is a life distribution, it is simply called a decreasing failure rate (DFR) distribution.

The set of EDFR distributions will be denoted by \mathcal{D} . An EDFR distribution F can have no jump on $(0, \infty)$ since R is concave on $(0, \infty)$, but can have a jump at 0. Also, notice that we consider δ_0 , the distribution degenerate at 0, to be an EDFR distribution. The hazard function of an EDFR distribution other than δ_0 is finite and concave on $(0, \infty)$. Consequently it is absolutely continuous on every closed interval of $(0, \infty)$ and has a right derivative which exists everywhere on $(0, \infty)$. In the remainder of this paper the failure rate function of an EDFR distribution other than δ_0 is always taken to be this right derivative and is, therefore, a de-

creasing function on $(0, \infty)$ with derivative a.e. less than or equal to zero. For convenience we shall define r to be identically zero when $F = \delta_0$.

Let L denote the class of extended life distributions, and $\sigma(L)$ denote the smallest σ -field of subsets of L such that the map $F \rightarrow F(t)$ from L into $[0, 1]$ is $\sigma(L)$ -measurable for all $t \in [0, \infty]$.

Definition 2.2. An extended life distribution F is a mixture of elements in a set $S \in L$ if $S \in \sigma(L)$ and there is a probability measure μ_F defined on $\sigma(L)$ such that

$$(3.1) \quad \mu_F(S) = 1$$

and

$$F(t) = \int G(t) \mu_F(dG) \text{ for all } t \in [0, \infty].$$

When (3.1) holds, μ is said to have support S .

Remark 2.3. It can be shown that $\sigma(L)$ is the Borel σ -field of L , when L is given the topology of weak convergence of probability measures. In particular, μ_F is a Borel measure.

We recall that an element x of a convex set K is an extreme point of K if $x = py + (1-p)z$ with $y, z \in K$ and $p \in (0, 1)$ implies that $y = z = x$.

We need Choquet's Theorem, stated below, to show that every EDFR life distribution can be represented as a mixture of extreme points of the EDFR class.

Choquet's Theorem (Phelps, 1966, pp. 19-20). Let K be a metrizable, compact, convex subset of a locally convex space X . Let $x_0 \in K$. Then there is a probability measure μ_{x_0} supported by the extreme points of K such that

$$L(x_0) = \int L(x) \mu_{x_0}(dx)$$

for all continuous linear functionals L defined on X .

In this paper we take X to be $M[0, \infty]$, the space of finite signed measures on $[0, \infty]$, with the topology of weak convergence, and K to be \mathcal{D} .

It is well known that with the above topology, $M[0, \infty]$ is locally convex and \mathcal{D} is metrizable, a convenient metric being the Lévy metric. An argument using Helly's Compactness Theorem and the fact that pointwise limits of concave functions are concave shows that \mathcal{D} is compact. The convexity of \mathcal{D} follows since the family of positive log-convex functions is closed under addition and multiplication (see Roberts and Varberg, 1973, Section 13).

To see that every EDFR can be represented as a mixture of extreme points of \mathcal{D} , we notice that $L: M[0, \infty] \rightarrow \mathbb{R}$ defined by $L(\nu) \equiv \int f(x)\nu(dx)$ is a continuous linear functional on $M[0, \infty]$ for each $f \in C[0, \infty]$. Hence the conclusion of Choquet's Theorem implies that for every EDFR distribution F there exists a probability measure, μ_F , supported by the extreme points of \mathcal{D} such that $\int f(x)F(dx) = \int [\int f(x)G(dx)]\mu_F(dG)$ for all $f \in C[0, \infty]$. Since $C[0, \infty]$ is a separating class of functions (see Breiman, 1968, p. 165) we have that $F(t) = \int G(t)\mu_F(dG)$ for all $t \in [0, \infty]$. Equivalently, by Remark 2.3, every EDFR distribution is a mixture of extreme points of \mathcal{D} . Further, by the definition of extreme point, we see that the set of extreme points of \mathcal{D} is the smallest set of EDFR distributions with the property that every EDFR may be represented as a mixture of its elements.

We identify the extreme points of \mathcal{D} in the next section.

3. The Extreme Points of the EDFR Class.

In this section we identify E , the set of extreme points of \mathcal{D} . In essence, we show that an EDFR distribution F is not an extreme point of \mathcal{D} if and only if either (i) $0 < F(0) < 1$, (ii) $0 < F(\infty) < 1$, or (iii) r' , the derivative of the failure rate function, is a.e. uniformly bounded away from 0 in some interval. More precisely, we show the following theorem.

Theorem 3.1. The EDFR distribution F is an extreme point of the set of EDFR distributions if and only if either (i) F is degenerate at 0, or at ∞ , or (ii) $F(0) = 0$, $F(\infty) = 1$, and $\{t: r'(t) \text{ exists, } a \leq t \leq b, \text{ and } r'(t) > -\delta\}$ has nonzero Lebesgue measure for all $0 < a < b$ and $\delta > 0$.

Let δ_∞ denote the distribution degenerate at ∞ .

Remark 3.2. The degree of smoothness of r for $F \in E$ characterizes certain types of extreme points. For example, let $F \in E \setminus \{\delta_0, \delta_\infty\}$. Then (i) r' is continuous if and only if F is an exponential distribution, and (ii) r' has a countable number of isolated discontinuities if and only if F is piecewise exponential, that is, R is piecewise linear. We remark that the EDFR piecewise exponentials are dense in \mathcal{D} . This example therefore shows that E is a dense subset of \mathcal{D} .

Two other types of interesting life distributions which are extreme points of \mathcal{D} are given below. These life distributions are extreme points of \mathcal{D} since their failure rates are decreasing functions with derivatives almost everywhere equal to 0. Notice that the failure rate function in (iii) below is discontinuous in a countable dense subset of $[0, \infty)$ while the failure rate function in (iv) below is continuous everywhere on $[0, \infty)$.

(iii) Let $\{a_j\}_{j=1}^{\infty}$ be a countable dense subset of $[0, \infty)$ and $\{b_j\}_{j=1}^{\infty}$ be a sequence of positive numbers such that $\sum b_j < \infty$ and $\sum a_j b_j = \infty$. For $t \geq 0$, let $r(t) \equiv \sum b_j I_{\{a_j \geq t\}}$. Then $F(t) \equiv 1 - e^{-\int_0^t r(u) du}$ is an extreme point of \mathcal{D} since $r' = 0$ a.e. on $[0, \infty)$, $F(0) = 0$, and $F(\infty-) = 1$.

(iv) Let $r(t)$ be positive, decreasing, continuous, and singular (with respect to Lebesgue measure). Further assume that $\int_0^{\infty} r(u) du = \infty$. Then $r' = 0$ a.e. on $[0, \infty)$, $F(0) = 0$, and $F(\infty-) = 1$. Consequently,

$F(t) = 1 - e^{-\int_0^t r(u) du}$ is an extreme point of \mathcal{D} .

In proving Theorem 3.1, we use the following notation. For each $F \in \mathcal{D}$ let $E_F \equiv \{t: r'_F \text{ exists at } t\}$. Denote the Lebesgue measure by m . Let $0 \equiv \{F \in \mathcal{D} \setminus \{\delta_0, \delta_{\infty}\}: m(t \in E_F: a \leq t \leq b \text{ and } r'(t) > -\delta) > 0 \text{ for all } 0 < a < b \text{ and all } \delta > 0\}$. Let $C \equiv \{F \in \mathcal{D}: F(0)=0 \text{ and } F(\infty-)=1\}$, i.e., C is the class of life distributions which place no mass at 0.

In this notation, Theorem 3.1 is equivalent to

$$(3.1) \quad E = (0 \cap C) \cup \{\delta_0, \delta_{\infty}\}.$$

We prove (3.1) by showing (i) $E \subset 0 \cup \{\delta_0, \delta_{\infty}\}$ (Lemma 3.3), (ii) $E \subset C \cup \{\delta_0, \delta_{\infty}\}$ (Lemma 3.4), and (iii) $E \supset (0 \cap C) \cup \{\delta_0, \delta_{\infty}\}$ (Lemma 3.5).

Lemma 3.3. $E \subset 0 \cup \{\delta_0, \delta_{\infty}\}$

Proof. It is enough to show that if $F \in \mathcal{D}$ and $F \notin 0 \cup \{\delta_0, \delta_{\infty}\}$, then $F \notin E$. To prove $F \notin E$, it suffices to show that there exist F_1 and $F_2 \in \mathcal{D}$ such that $(F_1 + F_2)/2 = F$.

Let $F \in \mathcal{D}$ and $F \notin 0 \cup \{\delta_0, \delta_{\infty}\}$. Then there exists an interval $[a, b]$, $(0 < a < b)$, and a value $\delta > 0$ such that $r'(t) < -\delta$ a.e. on $[a, b]$. Without loss of generality, we may assume that r is continuous at a and b . For each

positive integer n , let $\ell_n(x)$ be $= \frac{1}{n}(x-a)^3(b-x)^3$ for $a \leq x \leq b$ and 0 otherwise.

Choose n so that $\sup \ell_n < \ln 2$. Let ${}_nR_1 \equiv R - \ell_n$ and let ${}_nR_2 \equiv R + c(\ell_n)$, where $c(x) = -\ln(2-e^x)$ for $x < \ln 2$. We show that for n sufficiently large, say $n = n_0$, $F_i \equiv 1 - \exp\{{}_nR_i\}$ ($i=1,2$) are two different well-defined EDFR distributions such that $(F_1 + F_2)/2 = F$.

Since R is concave on $(0, \infty)$, it follows that r and r' exist a.e. on $[0, \infty)$. Further a.e. on $[0, \infty)$, we have that

$$\begin{aligned} {}_nR_1' &= r - \ell_n', \\ {}_nR_2' &= r + c'(\ell_n) \cdot \ell_n', \\ {}_nR_1'' &= r' - \ell_n'', \text{ and} \\ {}_nR_2'' &= r' + c'(\ell_n) \cdot \ell_n'' + c''(\ell_n)^2. \end{aligned}$$

Since $r'(t) < \delta$ a.e. on $[a, b]$, from the continuity of c' and c'' , it follows that for n sufficiently large, say $n = n_0$, ${}_nR_i(t) < -\delta/2$ a.e. on $[a, b]$ ($i=1,2$). Thus, writing R_i for ${}_nR_i$ ($i=1,2$), we have that R_i is strictly concave on $[a, b]$. We notice that R and R_i and also their derivatives agree off (a, b) . Thus R_i is an increasing function on $[0, \infty)$, is concave on $(0, \infty)$, and satisfies $R_i(\infty) \equiv \infty$. It follows that $F_i \equiv 1 - e^{-R_i}$ ($i=1,2$) are two different well-defined EDFR distributions.

We show that $(F_1 + F_2)/2 = F$. By the definitions of c , R_1 , and R_2 , we see that

$$R_2 = R - \ell_n (2 - e^{(R-R_1)}).$$

Hence

$$e^{-R_2} = e^{-R} (2 - e^{(R-R_1)}) = 2e^{-R} - e^{-R_1}.$$

Thus

$$\bar{F} = e^{-R} = (e^{-R_1} + e^{-R_2})/2 = (\bar{F}_1 + \bar{F}_2)/2. \quad ||$$

Lemma 3.4. $E \subset C \cup \{\delta_0, \delta_\infty\}$.

Proof. Let $F \in C \cup \{\delta_0, \delta_\infty\}$. Then either (i) $F(0) = \alpha$, where $0 < \alpha < 1$, or (ii) $F(\infty) = \beta$, where $0 < \beta < 1$. Assume (i) holds. Then

$F = \alpha \delta_0 + (1-\alpha) \left(\frac{F-F(0)}{1-F(0)} \right)$. Since $F_1 \equiv \delta_0$ and $F_2 \equiv \frac{F-F(0)}{1-F(0)}$ are both DFR

distributions, $F \notin E$. A similar argument shows $F \notin E$ when case (ii) holds. ||

Lemma 3.5. $(\mathcal{O} \cap C) \cup \{\delta_0, \delta_\infty\} \subset E$.

Proof. Let $F \in (\mathcal{O} \cap C) \cup \{\delta_0, \delta_\infty\}$. We show that $F \in E$. If $F = \delta_0$ or δ_∞ , then clearly $F \in E$. Assume then that $F \in \mathcal{O} \cap C$. Since \mathcal{D} is convex, it suffices to prove that

$$(3.2) \quad F = (F_1 + F_2)/2 \text{ with } F_1, F_2 \in \mathcal{D}$$

implies that

$$(3.3) \quad F_1 = F_2 = F.$$

Thus assume (3.2) holds. We show (3.3) follows.

For the distribution F_i ($i=1,2$), let r_i be the failure rate function and R_i be the hazard function. Let $t \in M \equiv \{t: r'_1 \text{ and } r'_2 \text{ exist at } t\}$. Since $m(M^c) = 0$ and $F \in \mathcal{O}$, it follows from the definition of \mathcal{O} that for each positive integer n ,

$$m([t, t+n^{-1}] \cap M \cap \{t: r'_i > -n^{-1}\}) > 0.$$

Hence we may choose a sequence $\{t_n\} \subset M$ such that $t_n \rightarrow t$ and $r'(t_n) \rightarrow 0$ as $n \rightarrow \infty$. By a subsequence argument we may also assume that the sequences $\{r'_1(t_n)\}$ and $\{r'_2(t_n)\}$ both have limits (possibly infinite). Now reversing the steps in the last paragraph of the proof of Lemma 3.3 we can show that $(R_2 - R)(t_n) = c((R - R_1)(t_n))$, where $c(x) \equiv -\ln(2 - e^x)$ for $x < \ln 2$. Differ-

entiating twice and taking limits, we get

$$(3.4) \quad \lim r_2'(t_n) = -c'((R-R_1)(t)) \cdot (\lim r_1'(t_n) + c''((R-R_1)(t)) \cdot (r(t) - r_1(t))^2,$$

since $r'(t_n) \rightarrow 0$ and $t_n \rightarrow t$ as $n \rightarrow \infty$ and R, R_1, r , and r_1 are continuous at t .

Now for $x < \ln 2$, $c'(x) = e^x/(2-e^x) > 0$ and $c''(x) = 2e^x/(2-e^x)^2 > 0$.

Hence, using the fact that $\lim r_1'(t_n) \leq 0$, we see that the expression on the right of (3.4) is nonnegative. Since the term on the left of (3.4) is non-positive, it follows that the expression on the right of (3.4) is equal to 0.

We conclude that $r(t) = r_1(t)$ for $t \in M$. Reversing the roles of R_1 and R_2 above, we get $r(t) = r_2(t)$ for $t \in M$. Thus

$$(3.5) \quad r = r_1 = r_2 \text{ a.e. on } [0, \infty).$$

Now $F(0) = 0$ implies that $F_1(0) = F_2(0) = F(0)$ and, consequently, that

$R(0) = R_1(0) = R_2(0)$. It follows by (3.5) that $R = R_1 = R_2$ or, equivalently, that $F = F_1 = F_2$. ||

4. Convex Extremal Subsets of \mathcal{D} .

In this section we identify the extreme points of two convex subclasses of \mathcal{D} , namely, (i) \mathcal{D}_p , the class of (proper) DFR distributions (Corollary 4.2), and (ii) $\mathcal{D}_C \equiv \{F \in \mathcal{D} : r_F \text{ is continuous on } (0, \infty)\}$ (Corollary 4.3). In addition in Theorem 4.5, we show that every DFR life distribution can be represented as a mixture of the extreme points of \mathcal{D}_p .

To identify the extreme points of \mathcal{D}_p and \mathcal{D}_C we need Lemma 3.1 below concerning extremal subsets. Recall that a subset K' of a convex set K is an extremal subset of K if $x = py + (1-p)z$ with $y, z \in K$ and $p \in (0,1)$ implies that y and z are in K' .

Let $\text{ext } K$ denote the set of extreme points of K .

Lemma 4.1. Let K' be a convex extremal subset of a convex set K . Then

$$\text{ext } K' = K' \cap \text{ext } K.$$

We omit the elementary proof of this lemma.

From Lemma 4.1 it follows that to identify the extreme points of \mathcal{D}_p and \mathcal{D}_C it is only necessary to show that these classes are convex extremal subsets of \mathcal{D} .

Since it is immediate that \mathcal{D}_p is a convex extremal subset of \mathcal{D} , we have the following corollary of Lemma 4.1.

Corollary 4.2. The DFR distribution F is an extreme point of the DFR class if and only if either (i) F is degenerate at 0, or (ii) $F(0) = 0$ and $\{t: r'(t) \text{ exists, } a \leq t \leq b, \text{ and } r'(t) > -\delta\}$ has nonzero Lebesgue measure for all $0 < a < b$ and $\delta > 0$.

To show that \mathcal{D}_C is an extremal subset of \mathcal{D} we need the following lemma. Let r, r_1 , and r_2 be the failure rates of F, F_1 , and F_2 .

Lemma 4.3. Let $F = pF_1 + (1-p)F_2$ for some $p \in (0,1)$ and $F_1, F_2 \in \mathcal{D}$. Let r be continuous at t . Then r_1 and r_2 are continuous.

Proof. Assume that either r_1 or r_2 is not continuous at t . We show that this contradicts the continuity of r at t .

Since r_1 and r_2 are decreasing and F_1 and F_2 are continuous on $(0, \infty)$, then

$$\begin{aligned} r(t+) &= [\bar{F}(t)]^{-1} [pr_1(t+)\bar{F}_1(t) + (1-p)r_2(t+)\bar{F}_2(t)] \\ &< [\bar{F}(t)]^{-1} [pr_1(t-)\bar{F}_1(t) + (1-p)r_2(t-)\bar{F}_2(t)] = r(t-), \end{aligned}$$

which contradicts the continuity of r at t . ||

It is clear from Lemma 4.3 that \mathcal{D}_C is an extremal subset of \mathcal{D} . Also a straightforward argument similar to that in the proof of Lemma 4.3 shows that \mathcal{D}_C is convex. Hence we obtain the following corollary of Lemma 4.1.

Corollary 4.4. The set of extreme points of \mathcal{D}_S is $E \cap \mathcal{D}_S$.

Remark 4.5. Let $\mathcal{D}_A \equiv \{F \in \mathcal{D} : r_F \text{ is absolutely continuous on every closed interval contained in } (0, \infty)\}$. It can be shown that \mathcal{D}_A is also a convex extremal subset of \mathcal{D} , and consequently that $\text{ext } \mathcal{D}_A = E \cap \mathcal{D}_A$.

We conclude this section with the following representation theorem for the DFR class.

Theorem 4.5. The life distribution F is DFR if and only if F may be represented as a mixture of distributions in $\text{ext } \mathcal{D}_p$. Further $\text{ext } \mathcal{D}_p$ is the smallest set with the property that every DFR may be represented as a mixture of its elements.

Proof. Let $F \in \mathcal{D}_p$. Then $F \in \mathcal{D}$ and, therefore, by the result stated

in the next to the last paragraph of Section 2, we have that
 $F(t) = \int G(t) \mu_F(dG)$ for all $t \in [0, \infty]$, where μ_F is a probability measure
 such that $\mu_F(E) = 1$. Since $F(\infty-) = 1$ by the Bounded Convergence Theorem,
 $1 = \int G(\infty-) \mu_F(dG)$. Hence $\mu_F(E \cap \mathcal{D}_p) = \mu_F\{G \in E : G(\infty-) = 1\} = 1$. Since
 $\text{ext } \mathcal{D}_p = E \cap \mathcal{D}_p$ by Corollary 4.2, the conclusion of the first part of
 Theorem 4.5 follows. The second part of Theorem 4.5 is an immediate
 consequence of the definition of an extreme point. ||

REFERENCES

1. Barlow, R. E. and Proschan, F. (1975). Statistical Theory of Reliability and Life Testing: Probability Models. Holt, Rinehart and Winston, Inc. New York.
2. Breiman, Leo (1968). Probability. Addison-Wesley Publishing Company, Reading Massachusetts.
3. Phelps, R. R. (1966). Lectures on Choquet's Theorem. Van Nostrand Co. Inc., Toronto.
4. Proschan, F. (1963). Theoretical explanation of observed decreasing failure rate. Technometrics 5 375-383.

UNCLASSIFIED

Classification of this Page

REPORT DOCUMENTATION PAGE

1. REPORT NUMBERS FSU No. M184-R AFCSR No. 78-5-R	2. GOVT. ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE The Extreme Points of the Set of Decreasing Failure Rate Distributions		5. TYPE OF REPORT & PERIOD COVERED 5 Technical Report
7. AUTHOR(s) Naftali A. Langberg Ramón V. León Frank Proschan		6. PERFORMING ORG. REPORT NUMBER FSU Statistics Report M484-R
8. CONTRACT OR GRANT NUMBER(s) AFOSR No. 78-3073 NSF No. MCS-7904698		9. PERFORMING ORGANIZATION NAME & ADDRESS The Florida State University Department of Statistics Tallahassee, Florida 32306
10. PROGRAM ELEMENT, PROJECT, TASK AREA AND WORK UNIT NOS.		11. CONTROLLING OFFICE NAME & ADDRESS U.S. Air Force Air Force Office of Scientific Research Bolling Air Force Base, D.C. 20332
12. REPORT DATE August, 1979		13. NUMBER OF PAGES 14
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS (of this report) Unclassified
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		16. DISTRIBUTION STATEMENT (of this Report) Approved for public release: distribution unlimited.
17. DISTRIBUTION STATEMENT (of the abstract, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS Decreasing failure rate, convex set, extreme points, Choquet's Theorem, mixture, hazard function, failure rate, reliability.		
20. ABSTRACT		

Since the class of extended decreasing failure rate (EDFR) life distributions (i.e., distributions with support in $[0, \infty]$) is compact and convex, it follows from Choquet's Theorem that every EDFR life distribution can be represented as a mixture of extreme points of the EDFR class. We identify the extreme points of this class and of the standard class of decreasing failure rate (DFR) life distributions. Further, we show that even though the convex class of DFR life distributions is not compact, every DFR life distribution can be represented as a mixture of extreme points of the DFR class.